



CrossMark
click for updates

Research

Cite this article: Smith SM, Fox RJ, Donelson JM, Head ML, Booth DJ. 2016 Predicting range-shift success potential for tropical marine fishes using external morphology. *Biol. Lett.* **12**: 20160505.
<http://dx.doi.org/10.1098/rsbl.2016.0505>

Received: 14 June 2016

Accepted: 6 September 2016

Subject Areas:

ecology, behaviour

Keywords:

biological invasion, climate change, coral reef fishes, morphological niche, tropicalization

Author for correspondence:

Shannen M. Smith

e-mail: shannen.m.smith@outlook.com

Electronic supplementary material is available online at <https://dx.doi.org/10.6084/m9.figshare.c.3472959>.

Global change biology

Predicting range-shift success potential for tropical marine fishes using external morphology

Shannen M. Smith¹, Rebecca J. Fox^{1,2}, Jennifer M. Donelson^{1,3}, Megan L. Head² and David J. Booth¹

¹School of Life Sciences, University of Technology Sydney, Ultimo, New South Wales 2007, Australia

²Division of Evolution, Ecology and Genetics, Research School of Biology, Australian National University, Canberra 2600, Australia

³ARC Centre of Excellence for Coral Reef Studies, James Cook University, Townsville, Queensland 4811, Australia

SMS, 0000-0003-1583-2785; MLH, 0000-0002-8123-7661

With global change accelerating the rate of species' range shifts, predicting which are most likely to establish viable populations in their new habitats is key to understanding how biological systems will respond. Annually, in Australia, tropical fish larvae from the Great Barrier Reef (GBR) are transported south via the East Australian Current (EAC), settling into temperate coastal habitats for the summer period, before experiencing near-100% mortality in winter. However, within 10 years, predicted winter ocean temperatures for the south-east coast of Australia will remain high enough for more of these so-called 'tropical vagrants' to survive over winter. We used a method of morphological niche analysis, previously shown to be an effective predictor of invasion success by fishes, to project which vagrants have the greatest likelihood of undergoing successful range shifts under these new climatic conditions. We find that species from the family of butterflyfishes (Chaetodontidae), and the moorish idol, *Zanclus cornutus*, are most likely to be able to exploit new niches within the ecosystem once physiological barriers to overwintering by tropical vagrant species are removed. Overall, the position of vagrants within the morphospace was strongly skewed, suggesting that impending competitive pressures may impact disproportionately on particular parts of the native community.

1. Introduction

Human-induced warming of the planet is driving shifts in the distributional ranges of many terrestrial [1] and marine [2] organisms. Predicting which species are most likely to be successful in establishing reproducing populations within their new habitat range is a critical step in forecasting the potential changes in biodiversity and ecosystem functioning that will occur as a result of continued global warming [3–5].

In the marine environment, poleward shifts in the distribution of tropical species are already resulting in the 'tropicalization' of temperate marine habitats [6]. The southeast coast of Australia lies at a known ocean warming hotspot [7,8] and is, therefore, likely to be particularly susceptible to the process of tropicalization. Every year tropical fish larvae are expatriated south from the Great Barrier Reef (GBR) via the East Australian Current (EAC), where they arrive and settle in recruitment pulses from January to May [9]. Currently, juvenile mortality reaches approximately 100% by July (Austral winter), driven by thermally induced reductions in physiological capacity. However, sea surface temperatures in this region are expected to rise by as much as 3°C by 2100, and coupled with the southward movement of the EAC [10], more of these tropical species are predicted to soon be capable overwintering

Table 1. Conditions of invader position within resident morphospace corresponding to probability of invasion success.

position within morphospace	probability of successful establishment in novel environment
falls well within convex hull of native community and occupies less morphological space than the nearest neighbours	low
positioned on the periphery of the native community, or occupies more morphological space than the nearest neighbours	medium
falls outside the original convex hull of the native community	high

at temperate latitudes. The winter ocean temperatures forecast to occur in the Sydney region by 2020–2030 [8] will be above the 17°C threshold currently estimated to be the constraint on physiological survival of tropical vagrant species [11], meaning that establishment of populations of tropical reef fishes in the waters off Sydney and the southeast coast of Australia will be possible within the next 15 years. Predicting which of these vagrant species are most likely to establish breeding populations, and therefore, compete with native species for food and habitat resources, is a critical issue for those concerned with managing these ecosystems into the future.

Many individual species-level traits have been proposed as predictors of geographical range-shift potential (e.g. high propagule production, generalist diet). However, the predictive power of the trait-based approach across multiple ecosystems and regions has been shown to be mixed, with some traits being identified as being important in certain contexts but not in others (reviewed in [12]). Recently, Azzurro *et al.* [13] demonstrated that external morphology alone can be an accurate predictor of a species' ability to successfully colonize a novel habitat. By using morphology as a proxy for a species' ecological niche, the method maps the existing resident community in terms of its morphological niche space and examines the position of potential invading species within this morphospace. Based on present-day abundances of species that have previously invaded the Mediterranean Sea via the Suez Canal, they showed that species which established successful populations were those that either lay outside, or at the margins of, the resident morphospace. Those that were morphologically similar to residents were unlikely to establish, presumably because the ecological niche was already filled [14,15]. Here we use this method to examine the position of seasonally invading tropical reef fish from the GBR within the morphospace of a native fish community of the southeast coast of Australia and to predict which vagrant species might be most likely to establish residency and shift their range as the physiological barriers to survival recede under global warming.

2. Material and methods

The morphological space of the native shoreline fish community of New South Wales (NSW), Australia was defined based on the species assemblage of a representative protected marine habitat (Cabbage Tree Bay, NSW, <https://ctbar.wordpress.com/>). A database of 110 species of bony fishes was compiled (electronic supplementary material, table S1) and an image of each species (left-hand side of adult individual) was sourced from online resources. Species' morphology was then characterized using

27 anatomical landmarks with ecological significance (following [13,16,17]; electronic supplementary material, table S1). Landmarking of images was done using tpsDig v. 1.40 [18], and scaled via a generalized least-squares procedure (generalized Procrustes) to eliminate the effect of isometric body-size variation. Relative warp analysis (tpsRelw v. 1.60) [19] was then used to obtain coordinates (relative warp scores) for each species within the reduced shape space [20]. The first two warps accounted for 47% of the observed morphological variation within the community and factor scores from these two warps were used to plot the position of individual species and define a morphospace of the native fish community in two dimensions. The convex hull of the native community (the smallest set of points enclosing all 110 species, see [21]) was then calculated using Qhull [22] and the resulting convex hull broken down into a series of Voronoi polygons to provide an estimate of the ecological space around each species [23]. For species at the periphery, Voronoi polygons are unbounded and use of the convex hull as a boundary, therefore, results in an underestimation of the area surrounding peripheral and hull points.

The analyses were repeated for a fish community containing the 110 native species plus the 11 tropical vagrant species most commonly observed as juveniles in the Sydney region over at least two consecutive summers in the past 15 years [24] (electronic supplementary material, table S1). Relative warp analysis yielded the coordinates of the vagrant species within the two-dimensional community morphospace, with Voronoi polygons giving conservative estimates of the amount of morphological space occupied by the vagrant species in relation to their 'closest' native neighbours (polygons do not allow for overlapping niches). Based on the position of the tropical species within the convex hull of the resident community (table 1), their distance from neighbouring species and Voronoi polygon area, the potential to establish successful populations in the future was classified (table 2). Since the receiving community could be precisely defined, and was based on a single specific habitat type, the position of vagrant species was considered in relation to the whole native fish community, rather than just in relation to pre-assigned guild members (see approach in Azzurro *et al.* [13]). This meant that consideration of vagrants' potential to establish a niche was not *a priori* constrained alongside species with an assumed similar functional role. Calculations of convex hulls, Voronoi polygons and nearest-neighbour Euclidean distances were carried out in MATLAB v. 9.0.0.

3. Results and discussion

Based on morphometric analysis, the species of tropical reef fishes with the greatest probability of exploiting an available niche within the existing resident assemblage are the butterflyfishes *Chaetodon auriga*, *C. flavirostris* and *C. vagabundus* and the zandlid *Zanclus cornutus* (figure 1 and table 2). The damselfishes *Abudefduf bengalensis*, *A. sexfasciatus* and

Table 2. Predicted likelihood of range expansion by tropical vagrant fish species into temperate marine communities of the southeast coast of Australia, based on morphology alone.

	species	Voronoi polygon area ($\times 10^{-3}$)	nearest-neighbour distance in morphospace	probability of establishing	polygon no. (figure 1b)
	<i>Zanclus cornutus</i>	5.80 ^a	0.155	high	1
	<i>Chaetodon flavirostris</i>	0.90 ^a	0.042	high	2
	<i>Chaetodon vagabundus</i>	1.00 ^a	0.031	high	4
	<i>Chaetodon auriga</i>	1.40 ^a	0.013	high	3
	<i>Abudefduf bengalensis</i>	1.90	0.009	medium	5
	<i>Abudefduf sexfasciatus</i>	1.00	0.027	medium	6
	<i>Acanthurus triostegus</i>	0.70	0.025	medium	9
	<i>Abudefduf vaigiensis</i>	0.40	0.012	medium	7
	<i>Abudefduf whitleyi</i>	0.22	0.020	medium–low	8
	<i>Pomacentrus coelestis</i>	0.27	0.007	low	11
	<i>Chromis nitida</i>	0.19	0.008	low	10

^aConservative estimate of niche space owing to constraint of convex hull boundary.

A. vaigiensis are predicted to have medium probability of establishing within the resident community, based on morphology and available niche space (figure 1 and table 2). Overall, the morphological positions of vagrant species were strongly skewed to the upper and lower left-hand corners of the convex hull.

Azzuro *et al.* [13] also found that species belonging to the family Chaetodontidae had a high invasion probability in the Mediterranean, but argued that habitat constraints (obligate association with coral of the species concerned) not accounted for in the methodology meant that they would remain rare. The butterflyfishes identified by our study as having strong range-shift potential actually show versatility in their habitat associations, even within reef environments. *Chaetodon auriga*, for example, can be observed in mangroves or rubble habitats, suggesting that they may be able to exploit the rocky shoreline habitats of southeast Australia, as predicted here. Additionally, they are not obligate coral feeders, exhibiting a more generalist diet including algae and invertebrates. The potential impact of climate change on the Australian temperate fish community could, therefore, be to expand niche opportunities for algal crevice feeders (*Z. cornutus*), algal and invertebrate feeders (*Chaetodon* spp.).

The most abundant tropical vagrant species currently found as juveniles along the NSW coast is the damselfish *A. vaigiensis*. In our analysis, this species is predicted to have a moderate likelihood of establishing abundant populations within the resident community, being located within, but in close proximity to, the boundary of the convex hull of the resident assemblage. Its position lies in close proximity to the resident species *Microcanthus strigatus* and its addition to the assemblage divides the niche space of *M. strigatus* into a smaller polygon, suggesting potential competition. Juvenile *A. vaigiensis* are commonly observed in close proximity to *M. strigatus*, using similar habitat following the summer recruitment event (S.M.S. 2015, personal observation) suggesting strong overlap in ecological niche and highlighting the predictive power of the morphological approach used here. Furthermore, the fact that the position of tropical vagrant species considered in this study is strongly skewed to the upper and lower left-hand corners of the convex hull suggests that, in ecological terms, the impending escalation in competition for niche space may impact disproportionately on particular parts of the native community.

Geometric morphometrics, while a useful predictive tool, is based on the assumption that competitive interactions




Figure 1. (a) Morphological position of species within the overall convex hull of a representative present-day assemblage of bony fishes from the southeast coast of Australia. Each dot represents a single species and outlines represent Voronoi polygons defining ecological niche space surrounding each species. (b) Convex hull of the present-day resident assemblage (dashed line) overlaid with assemblage including seasonally invading tropical vagrant fish species (peripheral solid line). Numbers inside Voronoi polygons refer to species listing in table 2.

between residents and invaders are driving the likelihood of population establishment success. For our study system, the model assumes that all commonly observed tropical fish that can be transported to Sydney via the EAC have the potential to interact with temperate species and have equal opportunity to use resources. The model then assesses this probability relative to competitive exclusion owing to morphological differences. It does not take into account environmental drivers of range-shift success. Model predictions are also critically related to the morphologies used to construct the morphospace. As Azzurro *et al.* [13] point out, morphological peculiarities not captured by the analysis can prove to be critical ecological novelties. For example, the lionfish (*Pterois* spp.), predicted only to have a moderate

chance of invasion success within the defined morphospace [13], has unfortunately become a highly successful invader in the Western Atlantic [25] and Mediterranean [26].

The value of the geometric morphometric approach, however, lies in the ability to generate hypotheses of range-expansion potential, which can be tested once ocean temperatures reach overwintering levels on a consistent basis. The growth of citizen science projects such as the Range Extension Database and Mapping Project (Redmap, <http://www.redmap.org.au>) and ongoing data collection by Reef Life Survey (<http://reeflifesurvey.com/>) will be an avenue to ground-truthing the predictions made here. For the temperate marine ecosystems of Australia's southeast coast, Redmap's database of year-on-year numbers of adult-sized tropical

vagrant fishes will provide a means by which the predictions made by the current study can be tested, once overwintering by all vagrant species becomes a physiological reality.

Data accessibility. Data deposited in Dryad: <http://dx.doi.org/10.5061/dryad.q0g60> [20].

Authors' contributions. S.M.S., R.J.F., J.M.D., D.J.B. designed the study. S.M.S., M.L.H. and R.J.F. collected data and performed the analyses, S.M.S., R.J.F., J.M.D. and D.J.B. interpreted the data. All authors

contributed to drafting of the manuscript, and subsequent edits made in response to reviewer comments. All authors approved the final version for publication and agree to be held accountable for the content therein.

Competing interests. The authors declare no competing interests.

Funding. This research was funded by the Ian Potter Foundation (J.M.D.) and UTS School of Life Sciences (S.M.S.).

Acknowledgement. We thank our two anonymous reviewers for their considered and helpful comments.

References

- Chen I-C, Hill JK, Ohlemüller R, Roy DB, Thomas CD. 2011 Rapid range shifts of species associated with high levels of climate warming. *Science* **333**, 1024–1026. (doi:10.1126/science.1206432)
- Perry AL, Low PJ, Ellis JR, Reynolds JD. 2005 Climate change and distribution shifts in marine fishes. *Science* **308**, 1912–1915. (doi:10.1126/science.1111322)
- Pearman PB, Guisan A, Broennimann O, Raddin CF. 2008 Niche dynamics in space and time. *Trends Ecol. Evol.* **23**, 149–158. (doi:10.1016/j.tree.2007.11.005)
- Ehrlén J, Morris WF. 2015 Predicting changes in the distribution and abundance of species under environmental change. *Ecol. Lett.* **18**, 303–314. (doi:10.1111/ele.12410)
- Parravicini V, Azzuro E, Kulbicki M, Belmaker J. 2015 Niche shift can impair the ability to predict invasion risk in the marine realm: an illustration using Mediterranean fish invaders. *Ecol. Lett.* **18**, 246–253. (doi:10.1111/ele.12401)
- Vergés A *et al.* 2014 The tropicalization of temperate marine ecosystems: climate-mediated changes in herbivory and community phase shifts. *Proc. R. Soc. B* **281**, 20140846. (doi:10.1098/rspb.2014.0846)
- Hobday AJ, Pecl GT. 2014 Identification of global marine hotspots: sentinels for change and vanguards for adaptation. *Rev. Fish Biol. Fish.* **24**, 415–425. (doi:10.1007/s11160-013-9326-6)
- Sen Gupta A, Brown JN, Jourdain NC, van Sebille E, Ganachaud A, Vergés A. 2015 Episodic and non-uniform shifts of thermal habitats in a warming ocean. *Deep Sea Res. II* **113**, 59–72. (doi:10.1016/j.dsr2.2013.12.002)
- Feary DA, Pratchett MS, Emslie M, Fowler AM, Figueira WF, Luiz OJ, Nakamura Y, Booth DJ. 2014 Latitudinal shifts in coral reef fishes: why some species do and others do not shift. *Fish Fish.* **15**, 593–615. (doi:10.1111/faf.12036)
- Ridgeway KR. 2007 Long-term trend and decadal variability of the southward penetration of the East Australian Current. *Geophys Res Lett.* **34**, L13613.
- Figueira WF, Booth DJ. 2010 Increasing ocean temperatures allow tropical fishes to survive overwinter in temperate waters. *Glob. Change Biol.* **16**, 506–516. (doi:10.1111/j.1365-2486.2009.01934.x)
- Sunday JM *et al.* 2015 Species traits and climate velocity explain geographic range shifts in an ocean-warming hotspot. *Ecol. Lett.* **18**, 944–953. (doi:10.1111/ele.12474)
- Azzuro E, Tuset VM, Lombarte A, Maynou F, Simberloff D, Rodríguez-Pérez A, Solé RV. 2014 External morphology explains the success of biological invasions. *Ecol. Lett.* **17**, 1455–1463.
- MacArthur R, Levins R. 1967 The limiting similarity, convergence, and divergence of coexisting species. *Am. Nat.* **101**, 377–385. (doi:10.1086/282505)
- Pianka ER. 1974 Niche overlap and diffuse competition. *Proc. Natl Acad. Sci. USA* **71**, 2141–2145. (doi:10.1073/pnas.71.5.2141)
- Farré M, Tuset VM, Maynou F, Recasens L, Lombarte A. 2013 Geometric morphology as an alternative for measuring the diversity of fish assemblages. *Ecol. Indicators* **29**, 159–166. (doi:10.1016/j.ecolind.2012.12.005)
- Recasens L, Lombarte A, Sánchez P. 2006 Teleostean fish composition and structure of an artificial reef and a natural rocky reef area in Catalonia (North Western Mediterranean). *Bull. Mar. Sci.* **78**, 71–82.
- Rohlf FJ. 2004 *Department of ecology and evolution*. Stony Brook, NY: State University of New York. tpsDig 1.4 and tpsUtil.
- Rohlf FJ. 2015 *Department of ecology and evolution*. Stony Brook, NY: State University of New York. tps Relative Warps v1.60.
- Smith S, Fox R, Donelson J, Head M, Booth D. 2016 Data from: Predicting range-shift success potential for tropical marine fishes using external morphology. Dryad Data Repository. (<http://dx.doi.org/10.5061/dryad.q0g60>)
- Cornwell WK, Schwiik DW, Ackerly DD. 2006 A trait-based test for habitat filtering: convex hull volume. *Ecology* **87**, 1465–1471. (doi:10.1890/0012-9658(2006)87[1465:ATTFHF]2.0.CO;2)
- Barber CB, Dobkin DP, Huhdanpaa HT. 1996 The Quickhull algorithm for convex hulls. *ACM Trans. Math. Softw.* **22**, 469–483. (doi:10.1145/235815.235821)
- Du F, Zhang XC, Sui Y, Shao M, Hu L, Shan L. 2012 The relationship between aboveground biomass and Voronoi area of coexisting species in an old-field community. *Pol. J. Ecol.* **60**, 479–489.
- Booth DJ, Figueira WF, Gregson MA, Brown L, Beretta G. 2007 Occurrence of tropical fishes in temperate southeastern Australia: role of the East Australian Current. *Estuar. Coast. Shelf Sci.* **72**, 102–114. (doi:10.1016/j.ecss.2006.10.003)
- Albins MA, Hixon MA. 2013 Worst case scenario: potential long-term effects of invasive predatory lionfish (*Pterois volitans*) on Atlantic and Caribbean coral-reef communities. *Environ. Biol. Fish.* **96**, 1151–1157. (doi:10.1007/s10641-011-9795-1)
- Kletou D, Hall-Spencer JM, Kleitou P. 2016 A lionfish (*Pterois miles*) invasion has begun in the Mediterranean Sea. *Mar. Biodiv. Rec.* **9**, 46. (doi:10.1186/s41200-016-0065-y)